



British Columbia Immunization Schedule as of February 2009

Age	Provincial Vaccines	Route	Comments	Other Vaccines
2 months	DTaP/IPV/HiB/HepB (INFANRIX hexa™)	IM		1 st Rotavirus vaccine (RotaTeq®) (three dose series).
	PneuC-7 (Prevnar®)	IM		
	Men-C conjugate (NeisVac-C™)	IM		
4 months	2 nd DTaP/IPV/HiB/HepB (INFANRIX hexa™)	IM		2 nd Rotavirus vaccine (RotaTeq®) (three dose series).
	2 nd PneuC-7 (Prevnar™)	IM		
6 months	3 rd DTaP/IPV/HiB/HepB (INFANRIX hexa™)	IM		3 rd Rotavirus vaccine (RotaTeq®) (three dose series). Influenza yearly for children >6 months (two half-doses in 1 st year).
	{PneuC-7 (Prevnar™)} for high risk only*			
12 months	2 nd Men-C conjugate (NeisVac-C™)	IM		Consider Hepatitis A vaccine (two dose series), especially if traveling.
	3 rd PneuC-7 (Prevnar™)	IM		
	MMR	SC		
	Varicella (Varivax® III)	SC	If susceptible	
18 months	4 th DTaP/IPV/HiB (PediaceL®)	IM		
	2 nd MMR	SC		
2 years				Consider Men-C/A/Y/W135 conjugate (Menactra®) anytime after age 2 years. If travelling, consider Dukoral® for Traveller's Diarrhea and a typhoid fever vaccine.
4-6 years	DTaP/IPV (Quadracel®)	IM		Consider second dose of Varicella (Varivax® III). Influenza - two doses needed if receiving for the first time under 9 years.
11 years (Gr. 6)	Men-C conjugate (NeisVac-C™)	IM		Consider Men-C/A/Y/W135 conjugate (Menactra®) instead.
	Hepatitis B (Recombivax®)(2 dose series)	IM	If not had as infant, use adult dosage.	
	HPV (Gardasil®) (3 dose series)	IM		
14 years (Gr. 9)	Tdap (Adacel®)	IM	Then every 10 years	HPV (Gardasil®) for girls Gr. 9 and older in 2007 as will miss start of provincial program.

* High risk = Sickle cell, asplenia (anatomic or functional), immunosuppression disorders, chronic heart/kidney/lung/liver disease, diabetes, CF, solid organ/stem cell transplants, cochlear implant recipients. These patients should also get Pneu-23 polysaccharide vaccine after age 2 years.

For further information and links to other references please see links at our website <http://www.bcpeds.ca>